



[Name of Document] DESCRIPTION

[Title of the Invention] METHOD FOR PREDICTING A DRUG TRANSPORT CAPABILITY BY ABCG2 POLYMORPHISMS

[Claims]

[Claim 1]

A method for predicting a drug transport capability of a mammalian cell comprising the steps of:

collecting a sample from a mammal,

determining a polymorphism of the nucleotide sequence of ABCG2 gene or a polymorphism of the amino acid sequence of ABCG2 polypeptide.

[Claim 2]

The method of claim 1, wherein said ABCG2 gene comprises a DNA consisting of the nucleotide sequence of SEQ ID NO:1, and said polymorphism of the nucleotide sequence is one or more of single nucleotide polymorphisms at positions selected from the group consisting of 34, 376 and 421 of SEQ ID NO:1.

[Claim 3]

The method of claim 2, wherein said single nucleotide polymorphism is selected from the group consisting of G34A, C376T and C421A.

[Claim 4]

The method of claim 2 or 3, wherein said polymorphism of the nucleotide sequence is determined by any one of methods selected from the group consisting of a direct sequencing method, TaqMan method, invader method, mass spectrometry, RCA method and DNA chip method.

[Claim 5]

The method of claim 1, wherein said ABCG2 polypeptide comprises an amino acid sequence of SEQ ID NO:2, and said polymorphism of the amino acid sequence is one or more of amino acid polymorphisms at positions selected from the group consisting of 12, 126, and 141 of SEQ ID NO:2.

[Claim 6]


The method of claim 5, wherein said amino acid polymorphism is an amino acid substitution of Val12Met or Gln141Lys, or a deletion of the amino acid sequence downstream from the position 126 of SEQ ID NO:2.

[Claim 7]

The method of claim 5 or 6, wherein said polymorphism of the amino acid sequence is determined by any one of methods selected from the group consisting of mass spectrometry, two-dimensional electrophoresis method, and protein chip method.

[Claim 8]

The method of any one of claims 1 to 7, wherein said drug is a compound represented by the following general formula (I):



(I)

[in the formula, X^1 and X^2 each independently represent a hydrogen atom, halogen atom or hydroxyl group, R represents a lower alkylamino group (wherein said lower alkylamino group may be substituted with a hydrogen atom, amino group, formylamino group, one to three hydroxyl group(s), a pyridyl group optionally having substituent(s), or thienyl group optionally having substituent(s)), and G represents a pentose group or hexose group (said pentose group or hexose group may be substituted with an amino group).]

[Claim 9]

A polynucleotide having a single nucleotide polymorphism(s) at one or more

position(s) selected from the group consisting of 34, 376 and 421 of SEQ ID NO:1, said polynucleotide comprising any one of the positions of said single nucleotide polymorphisms and consisting of at least 10 contiguous nucleotides of SEQ ID NO:1, or a complementary polynucleotide thereto.

[Claim 10]

The polynucleotide of claim 9, wherein said single nucleotide polymorphism is selected from the group consisting of G34A, C376T, C421A and single nucleotide polymorphisms complementary thereto.

[Claim 11]

A polynucleotide having one or more of the nucleotide polymorphisms in the polynucleotide sequence of SEQ ID NO:1, said polymorphism selected from the group consisting of nucleotide polymorphisms by which the translated amino acid at position 12 is methionine, one at position 126 is stop codon, and one at position 141 is lysine, and consisting of at least 10 contiguous nucleotides including one or more of nucleotides located at the site of said nucleotide polymorphisms, or a complementary polynucleotide thereto.

[Claim 12]

A pair of PCR primers which specifically hybridize to ABCG2 gene, and amplify a DNA fragment of a portion of said gene, wherein the amplified DNA fragment comprises a nucleotide(s) at position 34, 376 or 421 of SEQ ID NO:1

[Claim 13]

The pair of PCR primers of claim 12, said pair of PCR primers selected from the group consisting of SEQ ID NO:5 and 6; SEQ ID NO:9 and 10; and SEQ ID NO:11 and 12.

[Claim 14]

A polynucleotide which specifically hybridizes to ABCG2 gene, and which is capable of detecting a polymorphism(s) of ABCG2 gene at position 34, 376 or 421 of SEQ ID NO:1.

[Claim 15]

The polynucleotide of claim 14, which is capable of using in any one of methods selected from the group consisting of a direct sequencing method, TaqMan method, invader method, mass spectrometry, RCA method and DNA chip method.

[Claim 16]

A polypeptide having polymorphic mutation(s) to ABCG2 protein defined in the following (a) or (b), said polypeptide is a polymorphic mutant wherein one or both of amino acids at positions 12 and 141 of SEQ ID NO:2 are substituted with other amino acid(s), a polypeptide fragment comprising said substituted amino acid and at least 10 contiguous amino acid residues of said polymorphic mutant, or a polypeptide wherein the amino acid sequence downstream from the position 126 of SEQ ID NO:2 is deleted:

(a) a human ABCG2 polypeptide consisting of an amino acid sequence of SEQ ID No: 2,

(b) an isopolypeptide of (a) consisting of an amino acid sequence of SEQ ID NO:2, wherein one or several amino acids except for the amino acids at positions 12, 126 and 141, are deleted, substituted or added, and having a drug transport capability.

[Claim 17]

An antibody which specifically binds to the mutant ABCG2 polypeptide of claim 16.

[Claim 18]

A transformed cell which expresses an polypeptide having polymorphic mutation(s) to ABCG2 protein defined in the following (a) or (b), said polymorphic mutation(s) being one or both of amino acid substitutions Val12Met and Gln141Lys of the amino acid sequence of SEQ ID NO:2:

(a) a human ABCG2 polypeptide consisting of an amino acid sequence of SEQ ID NO:2,

(b) an isopolypeptide of (a) consisting of an amino acid sequence of SEQ ID NO:2, wherein one or several amino acids except for the amino acids at

positions 12, 126 and 141, are deleted, substituted or added, and having a drug transport capability.

[Claim 19]


A method for measuring a drug transport capability using the transformed cell of claim 18.

[Claim 20]

A method for diagnosing a drug sensitivity comprising the steps of: analyzing a biological sample from a subject, and determining the presence or absence of a polynucleotide of any one of claims 9 to 11, or a polypeptide of claim 16.

[Claim 21]

The method of claim 20, wherein the subject having said polynucleotide and/or said polypeptide is suggested to be sensitive to the compound represented by the following general formula (I):



(I)

[in the formula, X^1 and X^2 each independently represent a hydrogen atom, halogen atom or hydroxyl group, R represents a lower alkylamino group (wherein said lower alkylamino group may be substituted with a hydrogen atom, amino group, formylamino group, one to three hydroxyl group(s), a pyridyl group optionally having substituent(s), or thienyl group optionally having substituent(s)), and G represents a pentose group or hexose group (said pentose group or hexose group may be substituted

with an amino group).]

[Claim 22]

A kit for diagnosing a drug sensitivity comprising one or more of the following (a) to (f):

- (a) the polynucleotide of any one of claims 9 to 11,
- (b) the pair of primers of claim 12 or 13,
- (c) the polynucleotide of claim 14 or 15,
- (d) the polypeptide of claim 16,
- (e) the antibody of claim 17, and
- (f) the transformed cell of claim 18.

[Claim 23]

A computer system for analyzing data of ABCG2 polymorphism, comprising:

- (a) an input/output device,
- (b) a memory containing the polymorphism data, and
- (c) a central processing unit.

[Detailed Description of the Invention]

[0001]

[Field of the Invention]

The present invention relates to a polypeptide which excretes drugs such as cancer chemotherapeutic agents from a cell and to a gene coding therefor. More specifically, the present invention relates to a method for predicting a drug transport capability of a mammalian cell by determining a single nucleotide polymorphism(s) of ABCG2 gene and/or an amino acid polymorphism(s) of ABCG2 polypeptide and also to a polynucleotide, polypeptide, kit, and the like used for the method.

[0002]

[Prior Art]

Prediction of sensitivity to cancer chemotherapeutic drugs has been a subject in conventional cancer therapy by the cancer chemotherapeutic drugs. Anti-tumor activity of a chemotherapeutic drug shows a great difference depending on the type of

cancer cells and physical trait of each patient. A chemotherapeutic drug is highly effective for some patients while, a resistance to the drug is observed for other patients. In addition, although tumors are sensitive to chemotherapeutic drugs in early stages, they exhibit multidrug resistance afterward. In the conventional methods however, it is very difficult to judge whether a chemotherapeutic drug is effective to a specific patient.

[0003]

As a major cause for the difference of sensitivity to chemotherapeutic drugs, there is a difference in drug concentrations in cells due to the difference in drug excreting capability. In those cancer cells, each of the transporters which excrete the chemotherapeutic drugs out of the cell is a member of ABC transporter superfamily (ATP-binding cassette transporter superfamily) and is a group of molecules which is localized in cell membrane and transports the substrate utilizing an energy source such as ATP hydrolysis.

[0004]

As representative examples of the transporter, there have been reported P-glycoprotein (hereinafter, referred to as "P-gp") encoded on MDR1 gene and multidrug resistance-related proteins (hereinafter, referred to as "MRP") encoded on MRP subfamily genes such as MRP1, MRP2 and MRP3. P-gp is a molecular pump which was already known to be involved in multidrug resistance in multiple types tumor, while MRP is a transporter which was firstly found to be involved in multidrug resistance in lung cancer and, later, found to be expressed in other types of cancer as well (Cole, S. P. C. et al., Science, 258, 1650-1654 (1992) and Leslie, E. M. et al., Toxicology, 167, 3-23 (2001)).

[0005]

In recent years, new ABC family molecules have been found in succession and, besides P-gp and MRP, molecular pumps that are suggested to be involved in drug resistance are being clarified. As one of such molecules, there is a molecular pump called ABCG2 (BCRP/MXR/ABCP). With regard to this, there have been named and

reported ABCP as the gene which is expressed specifically in placenta (Allikmets, R. et al., Cancer Res. 58, 5337-5339 (1998)), BCRP as the gene obtained from a resistant cell line selected by adriamycin (Doyle, A. et al., Proc. Natl. Acad. Sci. U. S. A. 95, 15665-15670 (1998)) and MXR as the gene obtained from a resistant cell line selected by mitoxantrone (Miyake, K. et al., Cancer Res. 59, 8-13 (1999)). Among these three kinds of genes, mutations of 1 to 4 amino acid(s) derived from the nucleotide substitution between the respective genes were observed.

[0006]


From the analysis of the cell line which is produced by introducing and expressing the nucleotide sequence reported as BCRP into MCF-7 cell, expression of this gene was shown to give resistance to mitoxantrone and adriamycin. Thus, the gene has been notable for a novel factor of multidrug resistance (Doyle, A. et al., Proc. Natl. Acad. Sci. U. S. A., 95, 15665-15670 (1998) (WO 99/40110).

[0007]

Under such circumstances, the present applicant found that the excretion pump of indolocarbazole compounds is an ABCG2 gene of SEQ ID NO:1 (Komatani, H. et al., Cancer Research, 61, 2827-2832 (2001), WO 02/28894). In the gene reported as BCRP, the 482nd codon encodes threonine, while the ABCG2 gene of SEQ ID NO:1 was a new nucleotide sequence where the 482nd codon encodes arginine.

[0008]

The ABCG2 gene of SEQ ID NO:1 is a gene which confers a selective resistance on a cell to a compound of the following general formula (I) (hereinafter, referred to as "indolocarbazole compound"):



(I)

[in the formula, X1 and X2 each independently represents a hydrogen atom, halogen atom or hydroxyl group; R represents a lower alkylamino group (wherein said lower alkylamino group may be substituted with a hydrogen atom, amino group, formylamino group, one to three hydroxyl group(s), a pyridyl group optionally having substituent(s), or thienyl group optionally having substituent(s)); and G represents a pentose group or hexose group (said pentose group or hexose group may be substituted with an amino group).], more specifically, to the compound such as Compound A (wherein X1 is 1-hydroxyl group, X2 is 11-hydroxyl group, R is formylamino and G is β -D-glucopyranosyl group in the general formula (I)) and Compound B (wherein X1 is 2-hydroxyl group, X2 is 10-hydroxyl group, R is (1-hydroxymethyl-2-hydroxyl) ethylamino group and G is β -D-glucopyranosyl group in the general formula (I)).

[0009]

It has been shown by Northern blotting analysis that the ABCG2 gene of the SEQ ID NO:1, for example, is highly expressed in any cells which are resistant to both Compound A and Compound B (Yoshinari, T. et al., Cancer Res. 59, 4271-4275 (1999)) and that the accumulation of indolocarbazole compounds represented by Compound A, Compound B etc. into the cells is selectively suppressed by the gene. (Komatani, H. et al., Cancer Res. 61, 2827-2832 (2001); WO 02/28894). Accordingly,

analysis of the genetic polymorphisms affecting the activity or expression of ABCG2 comprising the ABCG2 gene of SEQ ID NO:1 is thought to be useful for the selection of anticancer drug used for the therapy. However, such a genetic polymorphism has not yet been known.

[0010]

[Problems to be solved by the Invention]

Under such circumstances, there has been a demand for the development of methods for diagnosis of excreting capability of a transporter gene product which excretes chemotherapeutic drugs out of the cells in each patient. For example, cancer chemotherapeutic drugs having an anthraquinone skeleton such as adriamycin, doxorubicin and mitoxantrone are not well effective to cells when the P-gp, MRP or BCRP is detected to be highly expressed therein.

[0011]

Although the indolocarbazole compounds are effective anti-cancer drugs regardless of the expression of the P-gp or MRP, their effect to cancer cells where ABCG2 is highly expressed is low.

[0012]

However, if the genetic polymorphism affecting the activity or the expression of ABCG2 can be previously detected, the detection may be useful for the selection of anti-cancer drugs in cancer therapy and for the selection of inhibitors of ABCG2 activity in combined cancer therapy.

[0013]

For example, the ABCG2 gene of SEQ ID NO:1 which is widely found is a gene giving an indolocarbazole compound-selective resistance on a cell while the ABCG2-Thr482 gene where the 482nd amino acid is modified to threonine gives a resistance to mitoxantrone and adriamycin in addition to indolocarbazole compounds and, therefore, a method for detecting the difference between those two genes is useful for the selection of anti-cancer drugs in cancer therapy.

[0014]

In addition, a detection of ABCG2 genetic polymorphism which lowers the activity of ABCG2 in advance, for example, is useful for finding the optimum dose of the indolocarbazole compound in cancer therapy.

[0015]

Accordingly, it is an object of the present invention to provide a polymorphism of ABCG2 polypeptide related to intracellular accumulation of indolocarbazole compounds and of a polynucleotide coding therefor. It is also an object of the present invention to provide a method for detecting the presence or absence of the polymorphism of ABCG2 polypeptide or polynucleotide coding therefor in the test sample derived from patients suffering from cancer, by using a nucleic acid which is specific to polymorphism of ABCG2-related gene or antibody to ABCG2 polypeptide. It is a still another object of the present invention to provide a method for an effective use of indolocarbazole compounds by detecting the presence or absence of the polymorphism of ABCG2 polypeptide or polynucleotide coding therefor.

[0016]

[Means to solve the Problems]

In order to solve the objects, the present inventors analyzed genomic DNA extracted from many human cancer cell lines and clinical samples and identified single nucleotide polymorphisms (SNPs) in the ABCG2 gene. It was found that those SNPs cause mutations such as an amino acid substitution and deletion at the specific sites of the ABCG2 polypeptide. Then, when cell lines expressing each of the specific mutant ABCG2 polypeptides were prepared and their resistance to drugs was tested, it was found that a drug transport capability of the mutant ABCG2 polypeptide greatly lowered as compared with that of wild type ABCG2 polypeptide. On the basis of such findings, the present invention has been accomplished.

[0017]

Accordingly, in a first aspect of the present invention, there is provided a method for predicting a drug transport capability of a mammalian cell comprising the

steps of collecting a sample from a mammal and determining at least a polymorphism of the nucleic acid sequence of ABCG2 gene or at least a polymorphism of the amino acid sequence of ABCG2 polypeptide.

[0018]

In a preferred embodiment of the present invention, the ABCG2 gene comprises a DNA consisting of the nucleotide sequence of SEQ ID NO:1 and the polymorphism of the nucleotide sequence is one or more of single nucleotide polymorphisms at positions selected from the group consisting of 34, 376 and 421 of SEQ ID NO:1. It is further preferred that the single nucleotide polymorphism is selected from the group consisting of G34A, C376T and C421A. Here, "G34A" means that the 34th guanine is substituted with adenine, "C376T" means that the 376th cytosine is substituted with thymine and "C421A" means that the 421st cytosine is substituted with adenine. Polymorphism of the nucleotide sequence can be determined by any one of methods selected from the group consisting of a direct sequencing method, TaqMan method, invader method, mass spectrometric method, RCA method and DNA chip method.

[0019]


In another preferred embodiment of the present invention, the ABCG2 polypeptide comprises a polypeptide consisting of an amino acid sequence of SEQ ID NO:2 and the polymorphism of the amino acid sequence is one or more of amino acid polymorphisms at positions selected from the group consisting of the 12, 126 and 141 of SEQ ID NO:2. It is preferred that the amino acid polymorphism is an amino acid substitution of Val12Met or Gln141Lys or deletion of the amino acid sequence downstream from the position 126 of SEQ ID NO:2. The polymorphism of the amino acid sequence can be determined by any of methods selected from the group consisting of mass spectrometric method, two-dimensional electrophoresis method and protein chip method.

[0020]

In a still preferred embodiment of the present invention, the aforementioned

drug is a compound represented by the following general formula (I) (hereinafter, referred to as "indolocarbazole compound").

[0021]



(I)

[In the formula, X^1 and X^2 each independently represent a hydrogen atom, halogen atom or hydroxyl group; R represents a lower alkylamino group (wherein said lower alkylamino group may be substituted with a hydrogen atom, amino group, formylamino group, one to three hydroxyl group(s), a pyridyl group optionally having substituent(s), or thienyl group optionally having substituent(s)); and G represents a pentose group or hexose group (said pentose group or hexose group may be substituted with an amino group).]

[0022]

In the second aspect of the present invention, there is provided a polynucleotide having a single nucleotide polymorphism(s) at one or more position(s) selected from the group consisting of 34, 376 and 421 of SEQ ID NO:1 wherein the polynucleotide comprises any one of the positions of the single nucleotide polymorphisms and consists of at least 10 contiguous nucleotides or a complementary polynucleotide thereto. In a preferred embodiment, the aforementioned single nucleotide polymorphism is selected from the group consisting of G34A, C376T, C421A and single nucleotide polymorphisms complementary thereto.

[0023]

In an embodiment, there is provided a polynucleotide having a nucleotide polymorphism(s) in the polynucleotide sequence of SEQ ID NO:1, wherein the polymorphism is one or more of nucleotide polymorphism(s) selected from the group consisting of nucleotide polymorphisms by which the translated amino acid at position 12 is methionine, one at position 126 is stop codon and one at position 141 is lysine, and comprising at least 10 contiguous nucleotides including one or more of nucleotide(s) located at the site of the nucleotide polymorphisms, or complementary sequence thereof.

[0024]

In the third aspect of the present invention, there is provided a pair of PCR primers which specifically hybridize to the ABCG2 gene and amplify a DNA fragment of a portion of the gene, wherein the amplified DNA fragment comprises a nucleotide at position 34, 376 or 421 of SEQ ID NO:1. In a preferred embodiment, the pair of PCR primers are any of the primer pairs selected from the group consisting of SEQ ID Nos. 5 and 6, SEQ ID Nos. 9 and 10 and SEQ ID Nos. 11 and 12.

[0025]

In the fourth aspect of the present invention, there is provided a polynucleotide which specifically hybridizes to ABCG2 gene and is capable of detecting the polymorphism of ABCG2 gene at position 34, 376 or 421 of SEQ ID NO:1. In a preferred embodiment, the aforementioned polynucleotide can be used in any of the methods selected from the group consisting of a direct sequencing method, TaqMan method, invader method, mass spectrometric method, RCA method and DNA chip method.

[0026]

In the fifth aspect of the present invention, there is provided a mutant ABCG2 polypeptide having polymorphic mutation(s) to either (a) a human ABCG2 polypeptide consisting of the amino acid sequence of SEQ ID NO:2 or (b) an isopolypeptide of (a) consisting of an amino acid sequence of SEQ ID NO:2 wherein one or several amino

acid(s) except for the amino acids at position 12, 126 and 141 are deleted, substituted or added and having a drug transport capability. And the said mutant ABCG2 polypeptide is a polypeptide where one or both of the amino acid(s) at positions 12 and 141 of SEQ ID NO:2 are substituted with other amino acid(s), or it is a polypeptide fragment comprising the substituted amino acid(s) and at least 10 contiguous amino acid residues of the mutant ABCG2 polypeptide above, or it is a polypeptide where the amino acid residues downstream from the position 126 of SEQ ID NO:2 are deleted.

[0027]

In the sixth aspect of the present invention, there is provided an antibody which specifically binds to the mutant ABCG2 polypeptide in the fifth aspect of the present invention.

[0028]

In the seventh aspect of the present invention, there is provided a transformed cell which expresses an ABCG2 polypeptide having one or both of amino acid substitutions Val12Met and Gln141Lys to the amino acid sequence of SEQ ID NO:2 defined as either (a) a human ABCG2 polypeptide consisting of an amino acid sequence of SEQ ID NO:2 or (b) an isopolypeptide of (a) consisting of an amino acid sequence of SEQ ID NO:2, wherein one or several amino acid(s) except for the amino acids at positions 12, 126 and 141, are deleted, substituted or added, and having a drug transport capability.

[0029]

In the eighth aspect of the present invention, there is provided a method for measuring a drug transport capability using the transformed cell in the seventh aspect.

[0030]

In the ninth aspect of the present invention, there is provided a method for diagnosing a drug sensitivity comprising the steps of collecting a sample from a subject and determining the presence or absence of the polynucleotide in the second aspect or the polypeptide in the fifth aspect. In a preferred embodiment, it is suggested that the subject having the polynucleotide and/or polypeptide is sensitive to

the indolocarbazole compound.

[0031]

In the tenth aspect of the present invention, there is provided a kit for the diagnosing a drug sensitivity comprising one or more of the polynucleotide in the second aspect, the pair of primers in the third aspect, the polynucleotide in the fourth aspect, the polypeptide in the fifth aspect, the antibody in the sixth aspect and the transformed cell in the seventh aspect.

[0032]

In the eleventh aspect of the present invention, there is provided a computer system for the analysis of ABCG2 polymorphism comprising (a) an input-output device(s), (b) a memory (storage medium) containing the polymorphism data and (c) a central processing unit.

[0033]

[Preferred Embodiments of the Invention]

(Definitions)

In the present specification, the following terms are defined as follows unless otherwise mentioned. "ABCG2" is a molecular pump which belongs to the ABC transporter superfamily and is a name of a polypeptide by which cancer chemotherapeutic drugs are excreted out of a cell or a gene coding therefor. The gene includes cDNA and genomic gene.

[0034]

The term "polymorphism" refers to the existence of more than one form of a gene, polypeptide or portion thereof. A portion of a gene, wherein there are at least two different forms, i.e., two different nucleotide sequences, is referred to as a "polymorphic region of a gene". A polymorphic region can be a single nucleotide pair, or can also be nucleotide pairs in some measure of length. The term "single nucleotide polymorphism (SNP)" refers to the replacement of one base by another base and, in human genome, it is presumed that an SNP is present in several hundreds to one thousand base pairs. Besides that, there exists some repetitive sequences having


a different repeated number among individuals in the site where a unit of two bases to several tens bases is repeatedly present, which are called VNTR (variable number of tandem repeats) and a microsatellite polymorphism. The SNPs have different functions depending upon the position where they are located, and some exist in the region which is translated to polypeptide resulting in substitution or deletion of amino acid sequence and affecting on the function of gene, some others exist in the region which controls the gene expression such as promoter or intron affecting on the expressed amount of gene, still some others exist in other region having nearly no influence on the gene expression.

[0035]

In the present specification, the term "mammalian cell" means tissues or cells that constitute animal body belonging to mammal or external cell cultures of such cells. The term "sample" means a sample containing polynucleotides derived from living organisms and includes living, dead or even archaeological sample collected from various tissues and cells. Specific examples are body fluid (blood, urine, saliva, and the like), skin, root of hair, mucous membrane, internal organs, placenta and cord blood.

[0036]

In the present specification, the term "drug" means a xenobiotic having a physiological activity including a cancer chemotherapeutic drug used for the purpose of treating cancer. It includes a synthetic compound, natural compound derived from plants or microorganisms and a semi-synthetic compound which is synthesized from the natural compound. Preferably, the "drug" means a compound represented by the following general formula (I) (hereinafter, referred to as "indolocarbazole compound"):



(I)

[In the formula, X¹ and X² each independently represent a hydrogen atom, halogen atom or hydroxyl group; R represents a lower alkylamino group (wherein said lower alkylamino group may be substituted with a hydrogen atom, amino group, formylamino group, one to three hydroxyl group(s), a pyridyl group optionally having substituent(s), or thienyl group optionally having substituent(s)); and G represents a pentose group or hexose group (said pentose group or hexose group may be substituted with an amino group).] More preferably, it means a compound of the general formula (I) [In the formula, X¹ and X² each independently represent a halogen atom or hydroxyl group; R represents a hydrogen atom, formylamino, or lower alkylamino wherein said lower alkylamino may be substituted with any one selected from the group consisting of one to three hydroxyl group(s), a pyridyl group optionally having substituent(s), and a thienyl group optionally having substituent(s); and G represents a hexose group which may be substituted with an amino group.] The production method and the like of the aforementioned indolocarbazole compounds have been disclosed in prior patent applications and registered patents (European patent publication 0528030 A1, U. S. Patent Nos. 5,591,842, 5,668,271, and 5,804,564, WO 95/30682, WO 96/04293, WO 98/07433 and JP Patent Kokai Publication No. JP-A-10-245390). Particularly with regard to the production methods of Compound A and Compound B, they are disclosed in JP Patent Kokai Publication No.

JP-A-6-128283 and WO 95/30682, respectively.

[0037]

In the present specification, the term "polynucleotide" generally refers to both polyribonucleotide and polydeoxyribonucleotide, which can be either a non-modified RNA or DNA and either a modified RNA or DNA. The examples thereof are DNA, cDNA, genomic DNA, mRNA, unprocessed RNA and fragments thereof. Although there is no particular limitation in its length, it is usually about 10 bases or longer. On the other hand, the term "oligonucleotide" refers to those which are relatively shorter than the "polynucleotide" in length, which is generally about 50 bases or less.

[0038]

In the present specification, the term "polypeptide" refers to a compound which is linked with a peptide bond(s) consisting of two or more amino acids and includes a relatively short-chain polypeptide called as a peptide or oligopeptide and a long-chain polypeptide called as a protein. The polypeptide may contain amino acid(s) which is other than the genetically coded 20 kinds of amino acids. It is also possible to contain modified amino acid. Such a modified amino acid(s) is produced *in vivo*, for example, by a posttranslational processing or by a chemical modification which is known among the persons skilled in the art. The modification can be take place at main chain of peptide bond, side chain of amino acid, amino terminal or carboxyl terminal and includes, for example, acetylation, acylation, ADP ribosylation, amidation, biotinylation, covalent bond with lipid or lipid derivative, formation of cross-linking bond, disulfide bond, addition of sugar chain, addition of GPI anchor, phosphorylation and prenylation.

[0039]

(Method for Predicting the Drug Transport Capability)

In an embodiment of the present invention, there is provided a method for predicting a drug transport capability of a mammalian cell, comprising the steps of collecting a sample from the mammal and determining at least a polymorphism of the nucleotide sequence of *ABCG2* gene or at least a polymorphism of the amino acid

sequence of ABCG2 polypeptide. Here, the ABCG2 gene comprises a human cDNA having a nucleotide sequence shown in SEQ ID NO:1 which is a gene giving a resistance on a cell to the indolocarbazole compound represented by Compound A. It further comprises a human isogene which hybridizes under a stringent condition to a DNA complementary to the nucleotide sequence of SEQ ID NO:1, and also encodes a polypeptide having a drug transport capability, as well as mammalian homologues thereto. The condition of "to hybridize under a stringent condition" is an experimental condition for hybridization which has been known among the persons skilled in the art. To be more specific, it means two nucleic acid fragments hybridize each other under a hybridization condition described in "Expression of cloned genes in *E. coli*" by J. Sambrook in 9.47 - 9.62 and 11.45 - 11.61 of "Molecular Cloning: A Laboratory Manual: 2nd edition (1989), Cold Spring Harbor Laboratory Press, New York, U. S. A."

[0040]

To be more specific, "under a stringent condition" means that, after hybridization at about 45°C in $6.0 \times \text{SSC}$, washing is conducted at 50°C in $2.0 \times \text{SSC}$. For the selection of stringency, salt concentration in the washing step may be selected, for example, from a low stringency of about $2.0 \times \text{SSC}$ at 50°C to a high stringency of about $0.2 \times \text{SSC}$ at 50°C. It is also possible that temperature for the washing step may be increased from a low stringency condition of room temperature or about 22°C to a high stringency condition of about 65°C. Incidentally, it is possible for persons skilled in the art to achieve a hybridization condition of the same stringency as the above condition by an appropriate selection of various conditions such as diluting ratio of SSC, concentration of formamide and temperature. Accordingly, the isogene includes various mutant genes which have been known already. For example, BCRP gene obtained from adriamycin-resistant cell line, ABCP gene which is specifically expressed in placenta and MXR gene obtained from resistant cell line selected by mitoxantrone are different from human ABCG2 gene of SEQ ID NO:1 in several nucleotide sequences but all of them are isogenes derived

from *ABCG2* gene of SEQ ID NO:1 and are included in "*ABCG2* gene" of the present invention.

[0041]

ABCG2 polypeptide is: (a) a human *ABCG2* polypeptide consisting of an amino acid sequence of SEQ ID NO:2; (b) an isopolypeptide to (a) consisting of an amino acid sequence of SEQ ID NO:2 wherein one or several amino acids except for the amino acid at positions 12, 126 and 141 are deleted, substituted or added and having a drug transport capability; or (c) a mammalian homologue to (a) or to (b). Here, the human *ABCG2* polypeptide consisting of the amino acid sequence of SEQ ID NO:2 is a polypeptide which gives a selective resistance on a cell to the indolocarbazole compound represented by Compound A. The isopolypeptide may have modifications of deletion, substitution or addition of one or several amino acid(s) in the amino acid sequence of SEQ ID NO:2 so far as a drug transport capability which is a function of the *ABCG2* polypeptide is remained, and numbers of modified amino acid in the functionally identical polypeptide are usually within 10% of total amino acids, preferably within 10 amino acids and, more preferably, modification numbers are within 3 amino acids (such as one amino acid).

[0042]

When the *ABCG2* genetic polymorphism and *ABCG2* polypeptide polymorphism are present in specific positions and those polymorphisms are present in more than a certain frequency in a specific population, the genetic significance of those polymorphisms becomes important. In a preferred embodiment of the present invention, specific SNPs as shown in Fig. 1 are disclosed. Fig. 1 shows schematically how the *ABCG2* polypeptide is present in cell membrane along with the SNP sites according to the present invention. The *ABCG2* polypeptide contains a leader sequence at its N-terminal necessary for localization to the cell membrane, followed by an ATP binding region (amino acids 61~270) and six transmembrane regions participating in the drug transportation. Fig. 1 shows four SNP sites which are mutations where the 34th guanine in SEQ ID No. 1 is substituted with adenine

(hereinafter, referred to as "G34A"), the 376th cytosine therein is substituted with thymine (hereinafter, referred to as "C376T"), the 421st cytosine therein is substituted with adenine (hereinafter, referred to as "C421A") and the 458th cytosine therein is substituted with thymine (hereinafter, referred to as "C458T"). As a result of those SNPs, in the amino acid sequence of the ABCG2 polypeptide, the 12th valine from the N-terminal is substituted with methionine (hereinafter, referred to as "Val12Met"), the 126th glutamine therefrom becomes a termination codon (hereinafter, referred to as "Gln126Term), the 141st glutamine therefrom is substituted with lysine (hereinafter, referred to as "Gln141Lys") and the 153rd threonine therefrom is substituted with methionine (hereinafter, referred to as "Thr153Met"). The mutation of Val12Met is present in a leader sequence necessary for localization of the ABCG2 polypeptide to cell membrane and the mutations of Gln141Lys and Thr153Met are present in an ABC (ATP-binding cassette) domain which is important for binding to ATP as transportation energy and, therefore, there is a strong possibility that those mutations affect on the drug transport capability of the ABCG2 polypeptide. It is apparent that a mutation of Gln126Term loses a drug transport activity since a complete ABCG2 polypeptide is not synthesized.

[0043]

Drug transport capability of those mutant ABCG2 polypeptides can be checked by preparing a transformant which expresses the mutant ABCG2 polypeptide by means of a recombinant DNA technique. As will be illustrated in detail hereinafter in the present specification, it is noted that, as a result of measurement of drug sensitivity using those transformed cells, the drug transport activity of the mutant ABCG2 polypeptides of the aforementioned Val12Met and Gln141Lys is significantly low as compared with that of the wild type ABCG2.

[0044]

Alternatively, it is also possible to test whether those mutations are related to the drug sensitivity or not by analyzing biological samples obtained from a group of the subjects having a high sensitivity to a specific drug and those obtained from a

normal group and then analyzing the statistical relationship with the polymorphism of the present invention (so-called case-control study). The statistical analysis can be carried out using a program, etc. being known among the persons skilled in the art.

[0045]

With regard to a method for determining the polymorphism of the aforementioned nucleic acid sequence, that may be carried out using various known arts which will be mentioned below for (1) determination of nucleotide sequence of a part of allele containing at least a polymorphic site, (2) detection by a probe (allele-specific probe) which specifically hybridizes to a polymorphic site, (3) measurement of molecular weight of gene fragment containing a polymorphic site, etc. For example, SNP can be directly detected from genomic DNA by a direct sequencing method. On the other hand, it is also possible to use the aforementioned means for identification of (1)~(3) after a specific genomic DNA region is amplified. Various methods for the DNA amplification are known to those skilled in the art and include, but are not limited to, cloning of a desired DNA fragment, polymerase chain reaction (PCR), ligase chain reaction (LCR), strand displacement amplification (SDA; Walker G., et al. *Proc. Natl. Acad. Sci. USA*, 89, 392-396 (1992)), transcription-based amplification (Kwoh, D. et al., *Proc. Natl. Acad. Sci. USA*, 86, 1173-1177 (1989)), self-sustained sequence replication (Guatelli, J., et al., *Proc. Natl. Acad. Sci. USA*, 87, 1874-1878 (1990)), the Q- β replicase system (Lizardi, P. et al., *Bio/Technology*, 6, 1197-1202 (1988)), nucleic acid sequence-based amplification (NASBA; Lewis, R., *Genetic Engineering News*, 12, 1 (1992)), the repair chain reaction (RCR), LAMP method (WO 00/28082), and the like.

[0046]

SNP of the amplified product can be determined by various methods, such as determination of the nucleotide sequence, measurement of molecular weight by MALDI-TOF mass spectrometry, etc. and analysis of restriction fragment length polymorphism (RFLP). The single strand conformation polymorphism (SSCP) detection technique is also such another method for separation based on an acrylamide

gel, etc., but non-denaturing conditions. It is also possible to carry out by a suitable capillary electrophoresis. This technique makes it possible to discriminate between different DNA fragments by their conformation (Orita, et al., *Proc. Natl. Acad. Sci. USA*, 86, (1989), *Cotton Mutat. Res.*, 285, 125-144 (1993), Hayashi, *Genet. Anal. Tech. Appl.* 9, 73-79 (1992)).

[0047]

TaqMan (trademark) method is a method for the detection of SNP using a fluorescent energy transfer phenomenon where hybridization of an allele-specific oligonucleotide to a template is performed simultaneously with PCR (cf. Livak, *et al.*, *PCR Methods and Application*, 4:357-362, 1995 and U. S. Patent No. 5,528,848). The allele-specific probe which is labeled with a fluorescent dye and a quencher is hybridized to a target site and PCR is performed using a primer which is designed to amplify the region including the site whereupon the hybridized probe is cleaved by 5'-nuclease activity of Taq polymerase as the elongation reaction from the primer proceeds. When the fluorescent dye is separated from the quencher, fluorescence is resulted and the template is amplified by the PCR whereupon fluorescent intensity is potentiated exponentially. When probes which are specific to two kinds of alleles are labeled with different fluorescent dyes, it is possible to discriminate homozygote from heterozygote by one assay.

[0048]

A variety of methods without amplification of DNA have been developed. For example, Invader method (trademark) is based on a special enzymatic reaction where two kinds of oligonucleotides (invader probe and allele probe) are used and a specific structure formed by those probes with a template DNA is recognized and cleaved and it is described, for example, in U. S. Patent Nos. 5,846,717, 5,614,402, 5,719,028, 5,541,311 and 5,843,669. In this method, the target nucleotide sequence is recognized by two different probes. The first probe is usually called an invader probe and is substantially complementary to the first site of the target nucleotide sequence. The second probe is called allele probe and its 3'-terminal side is substantially

complementary to the second site of the target nucleotide sequence while its 5'-terminal side contains a sequence called a tail or flap which is non-complementary to the template to form a single strand. When those probes hybridize to an adjacent region of the template, the 3'-terminal of the invader probe invades into an SNP site and this structure is cleaved by cleavase whereupon a flap is liberated. The liberated flap can be quantified when it is previously labeled. Preferably, in order to quantify the liberated flap, the third FRET (fluorescence resonance energy transfer) probe (including a sequence complementary to the flap and a self-complementary sequence) labeled with a fluorescent dye and a quencher may be used. The liberated flap forms a specific structure by binding to the FRET probe, and the part of fluorescent dye in the FRET probe is cleaved by cleavase whereupon fluorescence is generated. When two sets of flap-FRET probes are prepared and labeled with different fluorescent dyes, it is possible to discriminate each homozygote and heterozygote by one assay.

[0049]

MALDI-TOF mass spectrometry is a method which can process a large number of samples in a short time without fluorescence labeling of the primer. A primer adjacent to the SNP site is prepared and one base elongation reaction from the primer is performed using ddNTP and PCR-amplified sample DNA as a template. The ddNTP added to the elongation reaction product is discriminated by mass spectrometry.

[0050]

RCA (rolling circle amplification) is a method where a DNA amplifying means, in which a long complementary-stranded DNA is synthesized as DNA polymerase moves on a cyclic single-stranded template DNA, is applied to an SNP typing. Recognition of SNP (allele) is carried out by checking whether amplification is available by an RCA method. Namely, a single-stranded probe (padlock probe), which is able to give a ring form when annealed with genomic DNA, is hybridized to genomic DNA to conduct a chain reaction. When the terminal of the probe is made as a site of SNP, a connected ring is formed whereupon amplification by RCA takes

place if that site is matched while, when mismatched, no connection takes place giving no ring whereupon RCA amplification does not take place. Discriminating the two amplification reactions make it possible to determine SNP (Lizardi, P. M., et al., *Nat. Genet.*, 19, 225 (1998)).

[0051]

DNA chip method where PCR-amplified fluorescence-labeled cDNA or cRNA is hybridized to various oligonucleotide probes including polymorphism site using DNA chip arranged on a microarray is useful as a means for quick detection of many SNPs. There have been known a thing where oligonucleotide is synthesized on an array by an optical lithographic technique so that several thousands to several hundred thousands probes are arranged on a chip (manufactured by Affymetrix; cf. U. S. Patent Nos. 5,424,186, 5,744,101 and 6,040,138), and a method where a previously-prepared cDNA or oligonucleotide is fixed on glass by means of pin or ink jet system (cf. U. S. Patent No. 6,040,138).

[0052]

With regard to a method for determining the polymorphism of the aforementioned amino acid sequence, various methods have been known and examples thereof are a proteome analysis by a two-dimensional electrophoresis or microfluidics method (Vreeland, Wyatt N and Barron, Annelise E, *Current Opinion in Biotechnology*, Vol. 13, pages 87-94 (2002)), peptide mapping and amino acid sequence analysis using a mass spectrometric devices, amino acid sequence analysis by a protein sequencer and a method where interaction between polypeptide and ligand is detected using protein chips, etc.

[0053]

The two-dimensional electrophoresis is usually a method where isoelectric focusing is conducted in the first dimension while SDS-PAGE (sodium dodecyl sulfate-polyacrylamide gel electrophoresis) is conducted in the second dimension, and several thousands of proteins can be separated by one sheet of gel. In the isoelectric focusing, a carrier-ampholyte has been used already but, in recent years, immobilized

pH-gradient gel; IPG) strips have been put on the market and it is now possible to separate with a good reproducibility without causing a pH drift. In the SDS-PAGE, there are a continuous buffer system where one type of pH buffer is used and a discontinuous buffer system using buffers of plural pH values. It is also possible to use a low-BIS concentration gel electrophoresis, a concentration-gradient gel electrophoresis, a Tricine-SDS-PAGE, etc. depending upon the type of the protein to be separated. The separated protein can be usually quantified by staining with a dye such as Coomassie Blue. In a silver staining method, protein can be identified in a sensitivity of 20- to 100-fold as compared with the staining with Coomassie Blue. Alternatively, detection with a high sensitivity on a gel is possible using commercially available fluorescent dyes such as SYPRO Ruby and SYPRO Orange (Patton, W. F., *Electrophoresis*, 21, 1123-1144 (2000)). It is also possible to specifically detect an ABCG2 polypeptide by a western blotting method using an antibody to the ABCG2 polypeptide.

[0054]

Mass spectrometry is a technique by which mass (molecular weight) is precisely measured and, in recent years, this method makes it possible to measure precisely the molecular weights of the nucleic acids and proteins by making practical use of ionizing (highly hydrophilic) high-molecular substances having high polarity such as protein, etc. without decomposition. As one of such mass spectrometric methods, there has been known MALDI-TOF/MS (matrix-assisted laser desorption ionization time-of-flight/mass spectrometry). This is a method where, after mixing of a protein sample with a matrix which absorbs laser beam such as sinapinic acid (3,5-dimethoxy-4-hydroxycinnamic acid) followed by drying, a strong pulse laser is irradiated to conduct ionization of the protein sample by energy transfer from the matrix and molecular weight of the ion is analyzed by the difference in flying time of the molecular ion samples by the initial acceleration. In order to fragment the peptide in the inner area of the mass spectrometer and to obtain a structural information (such as amino acid sequence or amino acid composition) from analysis of the mass of the

fragment, a tandem mass spectrometry (MS/MS) where plural mass separating parts are connected is utilized and, for such a purpose, there are also used analyzers of a triple quadrupole type or a hybrid type using an electrospray ionization method and of an ion trap type, etc.

[0055]

Protein chip method is a technique which involves interaction of a sample with proteins, peptides, etc. placed on a substrate comprehensively and quickly and, with regard to ligands to be immobilized on the substrate, there have been developed peptides, antibodies, expressed proteins, etc.

[0056]

(Polynucleotide, Pair of primers and Kit)

In another embodiment of the present invention, there are provided a polynucleotide containing the *ABCG2* genetic polymorphism, a pair of primers for amplifying the DNA fragment containing the aforementioned polymorphisms, a polynucleotide for the detection of the aforementioned polymorphisms and a kit therefor.

[0057]

In an embodiment, the polynucleotide of this embodiment is a polynucleotide having a single nucleotide polymorphism(s) at one or more position(s) selected from the group consisting of 34, 376 and 421 of the SEQ ID NO:1 wherein the polynucleotide comprises any one of the positions of the aforementioned single nucleotide polymorphisms and consists of at least ten continuous nucleotides or complementary nucleotide thereto. Accordingly, a polynucleotide containing none of the single nucleotide polymorphisms at the aforementioned three positions or, in other words, a polynucleotide where the nucleotide sequences in the above three places are the same as those in SEQ ID NO:1 is not included in the present embodiment. Preferably, in those nucleotide sequences, the 34th guanine is substituted with adenine, the 376th cytosine is substituted with thymine or the 421st cytosine is substituted with adenine and, in addition to those, substitution in the nucleotide sequence where codon

of the 12th amino acid from N-terminal of the ABCG2 polypeptide is varied to methionine, the 141st codon thereof is varied to lysine or the 126th codon thereof is varied to termination codon is included as well.

[0058]

Those polynucleotides may be natural or synthetic compounds. For example, they may be manufactured by replication (duplication) of cDNA or genomic DNA within host cells using recombinant DNA techniques. Alternatively, they may be manufactured by synthesis *in vitro*. With regard to the synthetic method, it is possible to amplify the DNA by means of PCR or the like or to synthesize the DNA by chemical synthesis. For the persons skilled in the art, it is possible to introduce a site-specific mutation(s) into the ABCG2 gene of SEQ ID NO:1 by a known method to prepare the polynucleotide of the present embodiment. Examples of the method for introduction of site-specific mutations which is known to the persons skilled in the art are Kunkel method (Kunkel, T. A. et al., *Methods Enzymol.* 154, 367-382 (1987)), double primer method (Zoller, M. J and Smith, M., *Methods Enzymol.*, 154, 329-350 (1987)), cassette mutation method (Wells, et al., *Gene*, 34, 315-23 (1985)) and mega-primer method (Sarkar, G. and Sommer, S. S., *Biotechniques*, 8, 404-407 (1990)).

[0059]

Those polynucleotides may be used for the detection of the genetic polymorphisms concerning the present invention. They may be also used for the suppression of the gene expression as antisense DNAs.

[0060]

In another embodiment, there is provided a pair of PCR primers which specifically hybridize to the ABCG2 gene and amplify a DNA fragment of a portion of the gene wherein the amplified DNA fragment comprises a nucleotide at position 34, 376 or 421 of SEQ ID No. 1. The pair of primers of this embodiment are designed so as to be substantially complementary to each chain in specific regions of upper stream and downstream of the aforementioned polymorphic sites of ABCG2 gene. Although

each of those primers can be hybridized at the region which is apart in 25~2500 base pair, it is preferred that the size of the amplified product is 100~500 base pair so as to make determination of nucleotide sequence or analysis of molecular weight of the amplified product easy. More preferably, the size of the amplified product is 80~200 base pair. Although the length of those oligonucleotide primers may be within a range of 10~30 bases, there may be used an oligonucleotide primer having preferably 18~25 bases and, more preferably, an oligonucleotide primer having 20~22 bases as shown in SEQ ID Nos. 5 and 6, SEQ ID Nos. 9 and 10 and SEQ ID Nos. 11 and 12, respectively. Those primers may be labeled for making the detection of the amplified DNA fragment easy. With regard to the label, there may be used, for example, radioisotope, enzyme, fluorescent dye, streptoavidin, avidin, magnetic beads, antigen and antibody.

[0061]

In still another embodiment, there is provided a polynucleotide which specifically hybridizes to the *ABCG2* gene and which is capable of detecting a polymorphism of the *ABCG2* gene at positions 34, 376 or 421 of SEQ ID NO:1. With regard to a method for the detection, there are various methods and, for example, in conducting a detection by an invader method, there are provided an invader probe being designed to complementarily bind to the 3'-side of the template from the SNP site, and an allele probe containing a complementary sequence to the 5'-side of the template from the SNP site and having a sequence (flap) which is unrelated to the sequence of the template at the 5'-side thereof. The 3'-terminal of an invader probe which is a sequence of SNP site may be any base.

[0062]

A TaqMan probe is a polynucleotide which contains an SNP site and has a length of about 20 bases being complementary to a template. Its 5'-terminal is labeled with a fluorescent dye such as FAM or VIC while 3'-terminal is labeled with a quencher (optical quenching substance).

[0063]

In a padlock probe used for RCA method, each of its both ends comprises about 20 bases near the SNP on genome and two of them are linked by a specific sequence called a backbone.

[0064]

Preferably, the aforementioned detection is carried out by determining the nucleotide sequence using genomic DNA directly or using amplified DNA fragments. A sequence primer for the determination of a nucleotide sequence is designed so as to make it substantially complementary to an appropriate site in upper stream or downstream of the SNP site. Although the length of those sequence primers may be within a range of 10 to 30 bases, those having 18 to 25 bases may be preferably used and, more preferably, primers in an ordinary direction (sense primer) or primers in an inverted direction (antisense primer) as shown in SEQ ID Nos. 37, 38, 41, 42, 43 and 44 may be used. Those oligonucleotides may be chemically synthesized by various methods which are known among the persons skilled in the art. They may be also labeled for making the detection easy. With regard to a method of labeling, there may be used radioisotope, enzyme, fluorescent substance, streptoavidin, avidin, biotin, magnetic fine particles, antigen and antibody, etc.

[0065]

In another embodiment, there is provided a kit for the prediction and detection of drug transporting capability of mammals. The kit contains either or both of a pair of primers for amplifying the DNA fragments containing the ABCG2 polymorphism and a polynucleotide for detecting the polymorphism. It is possible that the target DNA is firstly amplified from the sample to be tested and genetic polymorphisms are determined using the amplified DNA. On the other hand, it is also possible to determine the polymorphism directly from genomic DNA without amplification reaction of DNA. With regard to such a method, Invader method may be exemplified. As optionally selected attachments, the kit may contain a reagent for extraction and for purification of DNA, reagent for PCR such as 10-fold concentrated buffer, heat-resistant DNA polymerase, four kinds of nucleotide triphosphates (dNTPs), etc.

[0066]

(Polypeptide)

In still another embodiment of the present invention, there is provided a polypeptide having a polymorphic mutation relevant to the present invention in the ABCG2 polypeptide or isopolypeptide thereof. The ABCG2 polypeptide is a polypeptide consisting of an amino acid sequence of SEQ ID NO:2 and the isopolypeptide thereof is a polypeptide consisting of an amino acid sequence where one or several amino acid(s) is/are deleted, substituted or added in the ABCG2 polypeptide and having a drug transport capability. Mutant polypeptides which have been known already such as BCRP, ABCP and MXR are also included therein. The polymorphism in this embodiment is substitution of either or both of the 12th and 141st amino acids in SEQ ID NO:2 with other amino acid(s) or deletion of an amino acid sequence which is downstream from the 126th of SEQ ID NO:2. The amino acid substitution may be anything so far as it is other than an amino acid residue shown in SEQ ID NO:2 but, preferably, the 12th and 141st amino acids in SEQ ID NO:2 are substituted with methionine and lysine, respectively.

[0067]

The present embodiment further includes polypeptide fragment of the ABCG2 polypeptide or isopolypeptide thereof where either or both of the 12th and 141st amino acid(s) of SEQ ID No. 2 is/are substituted with other amino acid(s). The polypeptide fragment comprises at least 10 contiguous amino acid residues, preferably 20 or more contiguous amino acid residues and, more preferably, it has the length of 30 or more amino acid residues. Those polypeptides or fragments thereof are useful for the preparation of an antibody to the polypeptide having polymorphic mutations.

[0068]

Such polypeptides having polymorphic mutation(s) can be manufactured by means of chemical synthesis and they further include natural polypeptides and those which are prepared as recombinant polypeptides utilizing genetic recombination techniques. The natural polypeptides may, for example, be an extracted and purified

polypeptide from tissues such as placenta where the human mutant ABCG2 polypeptide of this embodiment is thought to be expressed. On the other hand, the recombinant polypeptides can be prepared, as will be mentioned later, by cultivation of cells transformed by DNA coding for the human mutant ABCG2 polypeptide of this embodiment.

[0069]

The expressed or isolated polypeptide or the fragment thereof may be detected by known methods and it is possible to detect by, for example, Coomassie Blue staining, silver staining, western blotting method using an antibody specific to polypeptide having a polymorphic mutation, etc. In addition, those polypeptides may be purified by the methods which have been known already. Those methods include precipitation with ammonium sulfate, gel filtration chromatography, ion-exchange chromatography and affinity or immunochromatography.

[0070]

(Antibody)

In an embodiment of the present invention, an antibody which specifically binds to the mutant ABCG2 polypeptide having a polymorphism(s) of the present invention is provided. The antibody of this embodiment can be prepared according to the method known to the skilled person in the art (refer, for example, to "Shin Seikagaku Jikken Koza (New Experimental Course of Biochemistry) 1, Protein I, pages 389-406, Tokyo Kagaku Dojin"). The preparation of polyclonal antibody is performed, for example, as follows. To an immunocompetent animal such as rabbit, guinea pig, mouse and chicken is administered the appropriate dose of the mutant ABCG2 protein of the present invention or the partial peptide thereof. The administration may be accompanied by an adjuvant (FIA or FCA) which promotes the antibody production. It is generally administered every several weeks. Multiple immunizations can elevate the antibody titer. After the final immunization, antiserum is obtained by collecting blood from the immunized animal. The polyclonal antibody can be prepared from this antiserum by, for example, fractionation with ammonium

sulfate precipitation and/or anionic exchange chromatography and/or by affinity purification using Protein A and/or immobilized antigen. On the other hand, a monoclonal antibody is prepared, for example, as follows. The mutant ABCG2 polypeptide of the present invention or the partial peptide thereof is immunized to an immunocompetent animal as described above, and after the final immunization, spleen or lymph node is collected from the immunized animal. A hybridoma cell is prepared by the cell fusion of the antibody-producing cell which is contained in this spleen or lymph node and a myeloma cell using polyethylene glycol or the like. The aimed hybridoma is screened and cultivated and a monoclonal antibody can be prepared from the culture supernatant. Purification of the monoclonal antibody can be performed, for example, by fractionation with ammonium sulfate precipitation and/or anion exchange chromatography and/or by affinity purification using Protein A and/or immobilized antigen. The antibody thus prepared is used for an affinity purification of the mutant ABCG2 polypeptide of the present invention and may be used for detecting the amount of the expression of the mutant ABCG2 polypeptide of the present invention as well. The detection of the expressed amount of the mutant ABCG2 polypeptide of the present invention in a mammalian cell by the antibody makes it possible to determine the sensitivity of the mammalian cell to the compound represented by the formula (I). It is further possible that the detection of the mutant ABCG2 polypeptide of the present invention in a cancer cell or cancer patient by this antibody can be used for the pharmacogenomical therapy which determines the patient's constitution such as drug sensitivity for administration of optimum drug for the patient.

[0071]

(Transformed Cells and Method for Measuring the Drug Transport Capability Using the Transformed Cells)

The present invention further relates to a transformed cell which expresses an ABCG2 polypeptide having one or both of amino acid substitution(s) Val12Met and Gln141Lys of the amino acid sequence shown by SEQ ID NO:2 in: (a) a human

ABCG2 polypeptide consisting of an amino acid sequence of SEQ ID NO:2 or (b) an isopolypeptide of (a) consisting of an amino acid sequence of SEQ ID NO:2 wherein one or several amino acid(s) except for the amino acids at position(s) 12, 126 and 141 is/are deleted, substituted or added and having a drug transport capability. The term "transformed cell" refers to a cell where an exogenous DNA is incorporated into a host cell by a recombinant vector. The host cell may be either a prokaryotic cell or an eukaryotic cell and includes any cell which can be used for the object of the present invention, such as bacterium, yeast cell, insect cell or animal cell. To be more specific, it is possible to introduce the recombinant vector into the host cell by the following method whereupon the transformant is obtained. Transformation of *Escherichia coli* is carried out by the method of Hanahan (Hanahan, D., *J. Mol. Biol.* 166, 557-580 (1983)), the electroporation method (Dower, W. J., et al., *Nucl. Acid Res.* 16, 6127-6145 (1988)), and the like. Transformation of yeast is carried out, for example, by spheroplast method (Beach, D. and Nurse, P., *Nature*, 290, 140 (1981)), lithium acetate method (Okazaki, K., et al., *Nucleic Acids Res.*, 18, 6485-6489 (1990)), etc. Transformation of insect cell may be carried out by a method, for example, described in *Bio/Technology*, 6, 47-55 (1980). Introduction of recombinant DNA into mammalian cells is carried out by a calcium phosphate method (Graham, F. L. and van der Eb, A. J., *Virology*, 52, 456-467 (1973)), a DEAE-dextran method (Sussman, D. J. and Milman, G., *Mol. Cell Biol.*, 4, 1641-1643 (1984)), a lipofection method (Felgner, P. L. et al., *Proc. Natl. Acad. Sci. USA*, 84, 7413-7417 (1987)), an electric perforation method (Neumann, E., et al., *EMBO J.*, 1, 841-845 (1982)), etc. The transformed cells prepared as such may, for example, be used for a method for measuring the drug transporting activity, analysis of the drug excretion mechanism or screening of compound which regulates the drug transporting capability.

[0072]

(Diagnosing Method and Diagnosing Kit)

In a different embodiment of the present invention, there is provided a method for diagnosing a drug sensitivity of a subject by the detection of the polymorphisms of

the present invention or a kit therefor. The drug includes cancer chemotherapeutic drugs and it is clinically useful to diagnose the sensitivity therefor. For example, when a chemotherapeutic drug is administered to a specific patient suffering from cancer, responsiveness of the patient are different and there are big differences such as significantly effective, lowly effective and ineffective at all. This is because of a possibility that, since genetic background is different for each patient, activity of excreting the chemotherapeutic drug out of the cancer cell is greatly different. Accordingly, the diagnosis method of this embodiment is quite useful for deciding what type of chemotherapeutic drug or chemotherapeutic drug group is to be administered and/or for deciding the effective dose of chemotherapeutic drug or chemotherapeutic drug group. In a preferred embodiment, the subjects having polymorphisms shown in Table 3 are suggested to be sensitive to the indolocarbazole compound represented by the formula (I). Accordingly, a therapy by an effective dose of the said compound can be applied to the patient suffering from cancer for which the above compound is effective and a significant improvement in therapeutic effect as well as a big reduction in side effect can be expected.

[0073]

In another embodiment of the present invention, there is provided a kit for diagnosing a drug sensitivity containing one or more of the polynucleotide, pair of primers, polypeptide, antibody and transformed cell of the present invention. The diagnosing kit may contain an appropriate package for a safe storage of the constituting reagents and a package insert for illustrating the method of the present invention. It may further contain an appropriate buffer, nucleotide, polymerase such as heat-resistant polymerase and fluorescent substance for the detection.

[0074]

(Computer System)

In another embodiment of the present invention, there is provided a computer system where at least one SNP(s) of the *ABCG2* gene or at least one polymorphic mutant polypeptide sequence(s) concerning the *ABCG2* polypeptide is stored and

displayed. This computer system includes an input/output device, a central processing unit and a readable storage medium (memory) where the aforementioned polymorphic sequence data are stored. The above polymorphic sequence data include nucleotide sequence, genetic type and haplotype of the *ABCG2* gene in a subject population or amino acid sequence, spots by two-dimensional electrophoresis, mass spectrometric data, etc. of the *ABCG2* polypeptide. These data are processed by various programs and can be used for determination of genetic type, linkage disequilibrium analysis, etc. In a preferred embodiment, result of those analyses can be used for the prediction of drug sensitivity of mammalian cells.

[0075]

[Examples]

The present invention is explained in more detail by reference to the following examples which are results of identification of single nucleotide polymorphisms of the *ABCG2* gene using human genomic DNAs, and then preparation of cell lines expressing mutant *ABCG2* polypeptides to analyze the function thereof. However, these examples do not restrict the scope of the present invention.

[0076]

[Example 1] Identification of SNPs in Human *ABCG2* Gene

The present inventors firstly extracted genomic DNAs from 30 human cancer cell lines and also from human clinical samples of 149 persons (whites) and identified the SNPs by sequencing the *ABCG2* gene.

[0077]

The 30 cancer cell lines are A-427, DLD-1, NCI-H69, HeLa S3, PC-13, MKN-45, UM-UC-3, HCT116, PA-1, RT4, MKN1, SK-OV-3, MADH, KATOIII, U118, HS746, T24, MSTO-211H, OVCR3, Lu135, Lx-1, SCC25, Cal27, MKN-74, SCaBER, BxPC-3, Hela, J82, NCI-H 187 and ES-2. Genomic DNA was extracted from those cell lines with Trizol reagent (Gibco BRL). Human clinical samples were purchased from IMPATH-BCP Co. Nucleotide sequences of sixteen exons and peripheral introns of the *ABCG2* gene were determined by direct sequencing. Firstly,

sixteen exons were amplified from genomic DNA by PCR (LA Taq Takara) using each pair of primers shown in Table 1. Next, amplified DNA fragments were treated with ExoSAP-IT (USB corporation) to digest remaining primers and to remove unwanted dNTPs. Then, the DNA fragments were applied to cycle sequencing reaction with dye terminator method (Dynamic ET Dye Terminator Cycle Sequencing Kit; Amersham) using sense primers shown in Table 2. After the removal of dye-terminator by G-50 gel filtration column, the nucleotide sequences and SNPs were determined by capillary sequencer MegaBACE1000 (Molecular Dynamic). The identified SNPs were reconfirmed by sequencing using antisense primers shown in Table 2.

[0078]

Table 1

	Forward Primers	Reverse Primers
Exon 1	5'-GTGCCCACTCAAAAGGTT-3'	5'-TCCAGTCAAAGCTGTACTCTG-3'
Exon 2	5'-ATGTATTGTCACCTAGTGTG-3'	5'-AAAGTGTGAAGCCTTGAGCAGA-3'
Exon 3	5'-AACGGAGATGTTTCACAAGA-3'	5'-TACAATAAAGCCCCAAAACA-3'
Exon 4	5'-GAGGAAAAAGAATGGGAGAA-3'	5'-GTCTGCAAAGCCTGCTATAA-3'
Exon 5	5'-TTCCTTCACCTTCTTTTCC-3'	5'-CTTCCATAAAACTGGTCCCT-3'
Exon 6	5'-GAGGTGCTTTGTATCAGGCT-3'	5'-GATCAGGCCAGTAGGTCAAC-3'
Exon 7	5'-CTTGTAATACTTGCAGATTACCTG-3'	5'-TGTTCAAGTGACAGAATAAATGGCT-3'
Exon 8	5'-AAAGGGTAAATTACGTGGG-3'	5'-GCAAACAAACTGACGTTTTTC-3'
Exon 9	5'-AATGAAGGTGTTAGGGAAGC-3'	5'-CTGGCTGACACTTCTTTCAC-3'
Exon 10	5'-TCTCCCCAAAGCACAGATAACT-3'	5'-CATTTAAAAATAATTGGGCCAGGTG-3'
Exon 11	5'-CTAATTACCTTCCAAAGGGC-3'	5'-AAACCAGGCTGCTCTTTACT-3'
Exon 12	5'-GCTGGGTATTTTTCAAGGAT-3'	5'-AGAGAGTGCAAATGGACAG-3'
Exon 13	5'-TGCCTGTAGCTCTTCATCTC-3'	5'-ACGAGAGGGAACCAAAATAG-3'
Exon 14	5'-CTTTTTGGCAGCTTTAAATGATAGC-3'	5'-AATCTTTCTCCTTTACTAGGAGGTA-3'
Exon 15	5'-TTTACTTCTTTGTATTGGAAGCCA-3'	5'-TAGAGGATAAATCGATTGATAGGGA-3'
Exon 16	5'-ATCTGAAGGGTAATTATTAAGGC-3'	5'-TGTTCCAGAAATGGTGCAAGAATTC-3'

[0079]

Table 2

	Sense Primers	Antisense Primers
Exon 1	5'-GTGCCCACTCAAAAGGTT-3'	5'-CAAGAGTTTTTACCAACCCA-3'
Exon 2	5'-ATGTATTGTCACCTAGTGTG-3'	5'-GTGGCCCAATTATTTCACT-3'
Exon 3	5'-TAAGAGTTGGTTTGTGCTTG-3'	5'-AACATGGTCAACTGCTACAT-3'
Exon 4	5'-ATGTTTTGGGGCTTTATTG-3'	5'-TATTCCAGATTCTCCCTGC-3'
Exon 5	5'-CAGGCTTTCAGACATCTA-3'	5'-ATTGTTATGGAAAGCAACCA-3'
Exon 6	5'-GAGGTGCTTTGTATCAGGCT-3'	5'-CACCCTCATCACAGACATC-3'
Exon 7	5'-CTGTCCTAGAATCTGCATTT-3'	5'-AGCTGGTGCTACAAAAT-3'
Exon 8	5'-AAAGGGTAAATTACGTGGG-3'	5'-TCTGGTTGTTGCTTCCTACT-3'
Exon 9	5'-GTTAGGGAAGCATCCAAGA-3'	5'-AGGGAAGCTTTCAAAAGTA-3'
Exon 10	5'-TCTCCCCAAAGCACAGATAACT-3'	5'-TGGTGGTGGATGTCTGTAGT-3'
Exon 11	5'-CTAATTACCTTCCAAAGGGC-3'	5'-GCTCAGGATTTTCTTCCCTA-3'
Exon 12	5'-CTGGAAGTGTTCAGGAG-3'	5'-AGAGAGTGCAAATGGACAG-3'
Exon 13	5'-TGCCTGTAGCTCTTCATCTC-3'	5'-ATAAGGGCAAAGAGGAAAGT-3'
Exon 14	5'-TTTGTCTCTCCTTTAAACCG-3'	5'-AATCTTTCTCCTTTACTAGGAGGTA-3'
Exon 15	5'-TTTACTTCTTTGTATTGGAAGCCA-3'	5'-AAAAGGCCCAAAACAATAAG-3'
Exon 16-1	5'-ATCTGAAGGGTAATTATTAAGGC-3'	5'-CAGGAGTTTCCAGAATTCAA-3'
Exon 16-2	5'-TGTTGTTTTCTGTTCCCTTG-3'	5'-TGTTCCAGAAATGGTGCAAGAATTC-3'

[0080]

Results of the identified SNPs in the 30 human cancer cell lines and in the human clinical samples for 149 persons (whites) on the basis of the aforementioned determination of nucleotide sequences are shown in Table 3. In the column for domain in Table 3, ABC means ATP binding cassette, EC means extra cellular region, TM means transmembrane region and UTR means untranslated region. For example, when a mutation site is shown by counting the first adenine in the translation initiation codon as the 1st one, G34A was found in five cell lines (16.7%) in 30 kinds of cancer cell lines and in 29 persons (19.5%) in human clinical samples of 149 persons. Incidentally, the mutation where 10th adenine from the 5'-side of intron 3 is substituted with guanine is shown as "A+10G" and the mutation where 21st cytosine from the 3'-site of intron 13 is substituted with thymine is shown as "C-21T". Positions of some SNPs in Table 3 are shown in Fig. 1 together with a schematic structure of the ABCG2 polypeptide. Among those SNPs, G34A was present in a leader sequence which is important for localization of the ABCG2 polypeptide to a cell membrane and C421A was a mutation existing in an ATP binding cassette (ABC) region being important for binding to ATP which is transportation energy. Accordingly, those mutations have a high possibility of affecting the activity of the ABCG2 polypeptide. C376T is a mutation to termination codon existing in the ABC region and the fact that ABCG2 loses its activity is clear. Incidentally, each SNP for C496G, T623C, A1444G and G1445C is reported in NCBI SNP CLUSTER ID: rs 1061017, NCBI SNP CLUSTER ID: rs1061018, *Cancer Res.* 59, 8-13, 1999 and *Proc. Natl. Acad. Sci. USA*, 95, 15665-15670, 1998, respectively but they were not detected in the aforementioned cell lines and human-derived samples.

[0081]

Table 3

SNPs	Influence on Amino Acid Substitution	Existing Position	Domain	Frequency in 30 Cell Lines	Frequency in 149 Human Clinical Samples
G34C	Val12Met	Exon 2	Leader Sequence	5 (16.7%)	29 (19.5%)
A+10G	–	Intron 3	–	ND	25 (16.8%)
C369T	Tyr123Tyr	Exon 4	ABC	0	1 (0.7%)
C376T	Gln126Term	Exon 4	ABC	1 (3.3%)	0
C421A	Gln141Lys	Exon 5	ABC	6 (20%)	24 (16.1%)
C458T	Thr153Met	Exon 5	ABC	1 (3.3%)	0
C474T	Asp158Asp	Exon 5	ABC	0	1 (0.7%)
C496G	Gln166Glu	Exon 5	ABC	0	0
T623C	Phe208Ser	Exon 6	ABC	0	0
A+20G	–	Intron 11	–	ND	44 (29.5%)
A1444G	Arg482Gly	Exon 12	TM3	0	0
G1445C	Arg482Thr	Exon 12	TM3	0	0
C–21T	–	Intron 13	–	ND	36 (24.2%)
A1768T	Asp590Tyr	Exon 15	EC3	0	1 (0.7%)
G2237T	–	Exon 16	3'UTR	1 (3.3%)	0
G2393T	–	Exon 16	3'UTR	1 (3.3%)	0

[0082]

[Example 2] Preparation of Cell Lines Expressing mutated ABCG2

Among the polymorphic mutations identified in Example 1, two mutations – G34A and C421A – having a high possibility to affect the function of the ABCG2 polypeptide were prepared and introduced into animal cells as an endeavor to analyze their functions. Preparation of the mutated *ABCG2* genes was conducted by PCR and a point mutation was introduced. After confirming the introduction of the target mutations by sequencing, the mutated genes were cloned into *HindIII* and *XhoI* sites of an expression vector pcDNA3.1(+) and expression plasmid for each mutant was prepared. As a control, a plasmid expressing the wild type (WT) ABCG2 and the vector plasmid pcDNA3.1(+) alone without the *ABCG2* gene were used and those four kinds of expression plasmids were introduced to an animal cell (porcine kidney cell line) LLC-PK1 by lipofection method (Lipofectamine; Gibco BRL). Stable transfectants were selected with 1,500 $\mu\text{g/ml}$ of Geneticin (Gibco BRL) for two weeks

and cell lines were established. To determine the expressed level of ABCG2 in each cell line, total RNA was extracted from each transfectant cells and HeLa cells with Trizol (Gibco BRL). Seven microgram of the total RNA extracted from each clone of cell and full length ABCG2 cDNA (2.2 kb) probe labeled with ^{32}P were used for Northern hybridization. Several transfectants which expressed equal amount of ABCG2 mRNA were selected to eliminate the effect of expression level and the result is shown in Fig. 2. It is noted that, in Fig. 2, each of the transfectant cells of lanes 2-4 expressed ABCG2 mRNA in equal amount. Incidentally, GAPDH was used as an internal standard for mRNA expressed in each cell. It was noted that the lane 5 was a control clone which was transfected by vector alone and no ABCG2 mRNA was expressed.

[0083]

[Example 3] Evaluation of Resistance to Compound B

The transfectant cells which were selected in Example 2 and in which nearly equal amount of ABCG2 mRNA was expressed were incubated (cultivated) in a 199 medium containing 1 mM of L-glutamine, 50 units/ml of penicillin, 50 mg/ml of streptomycin and 10% by volume of fetal bovine serum. All of the incubations were carried out at 37°C under the humidified atmosphere containing 5% of carbon dioxide. The cytotoxicity of anticancer drugs was determined by sulforhodamine B dye-staining method and compared with each other. Specifically, four kinds of transformed cell clones were cultured at 37°C for 72 hours in a medium containing Compounds B or camptothecin of various concentrations, then fixed with trichloroacetic acid and stained for 30 minutes with 0.4% sulforhodamine B dissolved in 1% acetic acid solution. After unbound dye was removed by four washes with 1% acetic acid, polypeptide-bound dye was extracted with 10 mM unbuffered Tris base. Then, optical density of the extract was measured in a plate reader at 564 nm and 50% inhibitory concentration (IC_{50}) values for cell viability were determined. The results are shown in Table 4. In the cell line (1-58) expressing the wild type ABCG2, resistance to Compound B has been increased to an extent of 400-fold or more as

compared with the cell line (C4) which was transfected with vector alone. On the contrary, the resistance of the cell line (2-51) having a mutation of Val12Met in the leader sequence of ABCG2 or the cell line (3-28) having a mutation of Gln141Lys in the ABC region to Compound B increased to an extent of 7.7-fold and 48.2-fold, respectively, as compared with C4, however, the resistance as compared with that of wild type was about 1/10 or less. To camptothecin which is not a substrate for ABCG2, there was no significant difference in terms of the resistance among the cells. From these results, it was suggested that, the two kinds of mutant ABCG2 (Val12Met and Gln141Lys) obviously have a decreased capability of excreting Compound B which is a topoisomerase inhibitor out of the cell as compared with wild type ABCG2.

[0084]

Table 4 Influence of SNPs on Sensitivity to Compound B

Clone No.	C4	1-58	2-51	3-28
SNP Site	–	Wild Type	Leader Sequence (Val12Met)	ABC Domain (Gln141Lys)
Expression Level*	0	1.90	2.08	1.95
IC ₅₀ (μM Camptothecin)	0.0087	0.0213	0.0122	0.0268
IC ₅₀ (μM Compound B)	0.122	>50	0.94	5.88
Increased Rate	1.0	>409	7.7	48.2

*: Expression level of ABCG2 of each cell line was standardized to HeLa cell (= 1.0).

[0085]

[Effect of the Invention]

By using the method of the present invention, a drug transport capability of a mammalian cell can be predicted whereby sensitivity of a patient to various drugs such as anti-cancer drugs can be diagnosed and an indicator for the therapy can be obtained. In other words, as a result of selecting an anti-cancer drug in cancer therapy and, particularly, detecting a cancer cell(s) which is highly sensitive to indolocarbazole compounds, it is now possible to selectively apply the said compounds for the therapy. In addition, the optimum dose of the indolocarbazole compounds in the cancer therapy is found and, at the same time, side effect of the compounds is reduced whereby a highly effective method of using the indolocarbazole compounds is provided.

[0086]

[Sequence Listing]

SEQUENCE LISTING

<110> BANYU PHARMACEUTICAL CO., LTD

<120> A method for predicting a drug transporting capability
by ABCG2 polymorphisms

<130> P7194GN

<140>

<141>

<160> 68

<170> PatentIn Ver. 2.1

<210> 1

<211> 1968

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1).. (1965)

<400> 1

atg tct tcc agt aat gtc gaa gtt ttt atc cca gtg tca caa gga aac 48
Met Ser Ser Ser Asn Val Glu Val Phe Ile Pro Val Ser Gln Gly Asn
1 5 10 15

acc aat ggc ttc ccc gcg aca gct tcc aat gac ctg aag gca ttt act 96
Thr Asn Gly Phe Pro Ala Thr Ala Ser Asn Asp Leu Lys Ala Phe Thr
20 25 30

gaa gga gct gtg tta agt ttt cat aac atc tgc tat cga gta aaa ctg 144
Glu Gly Ala Val Leu Ser Phe His Asn Ile Cys Tyr Arg Val Lys Leu
35 40 45

aag agt ggc ttt cta cct tgt cga aaa cca gtt gag aaa gaa ata tta	192
Lys Ser Gly Phe Leu Pro Cys Arg Lys Pro Val Glu Lys Glu Ile Leu	
50 55 60	
tcg aat atc aat ggg atc atg aaa cct ggt ctc aac gcc atc ctg gga	240
Ser Asn Ile Asn Gly Ile Met Lys Pro Gly Leu Asn Ala Ile Leu Gly	
65 70 75 80	
ccc aca ggt gga ggc aaa tct tcg tta tta gat gtc tta gct gca agg	288
Pro Thr Gly Gly Gly Lys Ser Ser Leu Leu Asp Val Leu Ala Ala Arg	
85 90 95	
aaa gat cca agt gga tta tct gga gat gtt ctg ata aat gga gca ccg	336
Lys Asp Pro Ser Gly Leu Ser Gly Asp Val Leu Ile Asn Gly Ala Pro	
100 105 110	
cga cct gcc aat ttc aaa tgt aat tca ggt tac gtg gta caa gat gat	384
Arg Pro Ala Asn Phe Lys Cys Asn Ser Gly Tyr Val Val Gln Asp Asp	
115 120 125	
gtt gtg atg ggc act ctg acg gtg aga gaa aac tta cag ttc tca gca	432
Val Val Met Gly Thr Leu Thr Val Arg Glu Asn Leu Gln Phe Ser Ala	
130 135 140	
gct ctt cgg ctt gca aca act atg acg aat cat gaa aaa aac gaa cgg	480
Ala Leu Arg Leu Ala Thr Thr Met Thr Asn His Glu Lys Asn Glu Arg	
145 150 155 160	
att aac agg gtc att caa gag tta ggt ctg gat aaa gtg gca gac tcc	528
Ile Asn Arg Val Ile Gln Glu Leu Gly Leu Asp Lys Val Ala Asp Ser	
165 170 175	
aag gtt gga act cag ttt atc cgt ggt gtg tct gga gga gaa aga aaa	576
Lys Val Gly Thr Gln Phe Ile Arg Gly Val Ser Gly Gly Glu Arg Lys	
180 185 190	
agg act agt ata gga atg gag ctt atc act gat cct tcc atc ttg ttc	624
Arg Thr Ser Ile Gly Met Glu Leu Ile Thr Asp Pro Ser Ile Leu Phe	
195 200 205	

ttg gat gag cct aca act ggc tta gac tca agc aca gca aat gct gtc	672
Leu Asp Glu Pro Thr Thr Gly Leu Asp Ser Ser Thr Ala Asn Ala Val	
210 215 220	
ctt ttg ctc ctg aaa agg atg tct aag cag gga cga aca atc atc ttc	720
Leu Leu Leu Leu Lys Arg Met Ser Lys Gln Gly Arg Thr Ile Ile Phe	
225 230 235 240	
tcc att cat cag cct cga tat tcc atc ttc aag ttg ttt gat agc ctc	768
Ser Ile His Gln Pro Arg Tyr Ser Ile Phe Lys Leu Phe Asp Ser Leu	
245 250 255	
acc tta ttg gcc tca gga aga ctt atg ttc cac ggg cct gct cag gag	816
Thr Leu Leu Ala Ser Gly Arg Leu Met Phe His Gly Pro Ala Gln Glu	
260 265 270	
gcc ttg gga tac ttt gaa tca gct ggt tat cac tgt gag gcc tat aat	864
Ala Leu Gly Tyr Phe Glu Ser Ala Gly Tyr His Cys Glu Ala Tyr Asn	
275 280 285	
aac cct gca gac ttc ttc ttg gac atc att aat gga gat tcc act gct	912
Asn Pro Ala Asp Phe Phe Leu Asp Ile Ile Asn Gly Asp Ser Thr Ala	
290 295 300	
gtg gca tta aac aga gaa gaa gac ttt aaa gcc aca gag atc ata gag	960
Val Ala Leu Asn Arg Glu Glu Asp Phe Lys Ala Thr Glu Ile Ile Glu	
305 310 315 320	
cct tcc aag cag gat aag cca ctc ata gaa aaa tta gcg gag att tat	1008
Pro Ser Lys Gln Asp Lys Pro Leu Ile Glu Lys Leu Ala Glu Ile Tyr	
325 330 335	
gtc aac tcc tcc ttc tac aaa gag aca aaa gct gaa tta cat caa ctt	1056
Val Asn Ser Ser Phe Tyr Lys Glu Thr Lys Ala Glu Leu His Gln Leu	
340 345 350	
tcc ggg ggt gag aag aag aag aag atc aca gtc ttc aag gag atc agc	1104
Ser Gly Gly Glu Lys Lys Lys Lys Ile Thr Val Phe Lys Glu Ile Ser	
355 360 365	

tac acc acc tcc ttc tgt cat caa ctc aga tgg gtt tcc aag cgt tca 1152
 Tyr Thr Thr Ser Phe Cys His Gln Leu Arg Trp Val Ser Lys Arg Ser
 370 375 380

ttc aaa aac ttg ctg ggt aat ccc cag gcc tct ata gct cag atc att 1200
 Phe Lys Asn Leu Leu Gly Asn Pro Gln Ala Ser Ile Ala Gln Ile Ile
 385 390 395 400

gtc aca gtc gta ctg gga ctg gtt ata ggt gcc att tac ttt ggg cta 1248
 Val Thr Val Val Leu Gly Leu Val Ile Gly Ala Ile Tyr Phe Gly Leu
 405 410 415

aaa aat gat tct act gga atc cag aac aga gct ggg gtt ctc ttc ttc 1296
 Lys Asn Asp Ser Thr Gly Ile Gln Asn Arg Ala Gly Val Leu Phe Phe
 420 425 430

ctg acg acc aac cag tgt ttc agc agt gtt tca gcc gtg gaa ctc ttt 1344
 Leu Thr Thr Asn Gln Cys Phe Ser Ser Val Ser Ala Val Glu Leu Phe
 435 440 445

gtg gta gag aag aag ctc ttc ata cat gaa tac atc agc gga tac tac 1392
 Val Val Glu Lys Lys Leu Phe Ile His Glu Tyr Ile Ser Gly Tyr Tyr
 450 455 460

aga gtg tca tct tat ttc ctt gga aaa ctg tta tct gat tta tta ccc 1440
 Arg Val Ser Ser Tyr Phe Leu Gly Lys Leu Leu Ser Asp Leu Leu Pro
 465 470 475 480

atg agg atg tta cca agt att ata ttt acc tgt ata gtg tac ttc atg 1488
 Met Arg Met Leu Pro Ser Ile Ile Phe Thr Cys Ile Val Tyr Phe Met
 485 490 495

tta gga ttg aag cca aag gca gat gcc ttc ttc gtt atg atg ttt acc 1536
 Leu Gly Leu Lys Pro Lys Ala Asp Ala Phe Phe Val Met Met Phe Thr
 500 505 510

ctt atg atg gtg gct tat tca gcc agt tcc atg gca ctg gcc ata gca 1584
 Leu Met Met Val Ala Tyr Ser Ala Ser Ser Met Ala Leu Ala Ile Ala
 515 520 525

gca ggt cag agt gtg gtt tct gta gca aca ctt ctc atg acc atc tgt 1632
 Ala Gly Gln Ser Val Val Ser Val Ala Thr Leu Leu Met Thr Ile Cys
 530 535 540

ttt gtg ttt atg atg att ttt tca ggt ctg ttg gtc aat ctc aca acc 1680
 Phe Val Phe Met Met Ile Phe Ser Gly Leu Leu Val Asn Leu Thr Thr
 545 550 555 560

att gca tct tgg ctg tca tgg ctt cag tac ttc agc att cca cga tat 1728
 Ile Ala Ser Trp Leu Ser Trp Leu Gln Tyr Phe Ser Ile Pro Arg Tyr
 565 570 575

gga ttt acg gct ttg cag cat aat gaa ttt ttg gga caa aac ttc tgc 1776
 Gly Phe Thr Ala Leu Gln His Asn Glu Phe Leu Gly Gln Asn Phe Cys
 580 585 590

cca gga ctc aat gca aca gga aac aat cct tgt aac tat gca aca tgt 1824
 Pro Gly Leu Asn Ala Thr Gly Asn Asn Pro Cys Asn Tyr Ala Thr Cys
 595 600 605

act ggc gaa gaa tat ttg gta aag cag ggc atc gat ctc tca ccc tgg 1872
 Thr Gly Glu Glu Tyr Leu Val Lys Gln Gly Ile Asp Leu Ser Pro Trp
 610 615 620

ggc ttg tgg aag aat cac gtg gcc ttg gct tgt atg att gtt att ttc 1920
 Gly Leu Trp Lys Asn His Val Ala Leu Ala Cys Met Ile Val Ile Phe
 625 630 635 640

ctc aca att gcc tac ctg aaa ttg tta ttt ctt aaa aaa tat tct taa 1968
 Leu Thr Ile Ala Tyr Leu Lys Leu Leu Phe Leu Lys Lys Tyr Ser
 645 650 655

<210> 2

<211> 655

<212> PRT

<213> Homo sapiens

<400> 2

Met Ser Ser Ser Asn Val Glu Val Phe Ile Pro Val Ser Gln Gly Asn

1 5 10 15
Thr Asn Gly Phe Pro Ala Thr Ala Ser Asn Asp Leu Lys Ala Phe Thr
20 25 30
Glu Gly Ala Val Leu Ser Phe His Asn Ile Cys Tyr Arg Val Lys Leu
35 40 45
Lys Ser Gly Phe Leu Pro Cys Arg Lys Pro Val Glu Lys Glu Ile Leu
50 55 60
Ser Asn Ile Asn Gly Ile Met Lys Pro Gly Leu Asn Ala Ile Leu Gly
65 70 75 80
Pro Thr Gly Gly Gly Lys Ser Ser Leu Leu Asp Val Leu Ala Ala Arg
85 90 95
Lys Asp Pro Ser Gly Leu Ser Gly Asp Val Leu Ile Asn Gly Ala Pro
100 105 110
Arg Pro Ala Asn Phe Lys Cys Asn Ser Gly Tyr Val Val Gln Asp Asp
115 120 125
Val Val Met Gly Thr Leu Thr Val Arg Glu Asn Leu Gln Phe Ser Ala
130 135 140
Ala Leu Arg Leu Ala Thr Thr Met Thr Asn His Glu Lys Asn Glu Arg
145 150 155 160
Ile Asn Arg Val Ile Gln Glu Leu Gly Leu Asp Lys Val Ala Asp Ser
165 170 175
Lys Val Gly Thr Gln Phe Ile Arg Gly Val Ser Gly Gly Glu Arg Lys
180 185 190
Arg Thr Ser Ile Gly Met Glu Leu Ile Thr Asp Pro Ser Ile Leu Phe
195 200 205
Leu Asp Glu Pro Thr Thr Gly Leu Asp Ser Ser Thr Ala Asn Ala Val
210 215 220

Leu Leu Leu Leu Lys Arg Met Ser Lys Gln Gly Arg Thr Ile Ile Phe
225 230 235 240

Ser Ile His Gln Pro Arg Tyr Ser Ile Phe Lys Leu Phe Asp Ser Leu
245 250 255

Thr Leu Leu Ala Ser Gly Arg Leu Met Phe His Gly Pro Ala Gln Glu
260 265 270

Ala Leu Gly Tyr Phe Glu Ser Ala Gly Tyr His Cys Glu Ala Tyr Asn
275 280 285

Asn Pro Ala Asp Phe Phe Leu Asp Ile Ile Asn Gly Asp Ser Thr Ala
290 295 300

Val Ala Leu Asn Arg Glu Glu Asp Phe Lys Ala Thr Glu Ile Ile Glu
305 310 315 320

Pro Ser Lys Gln Asp Lys Pro Leu Ile Glu Lys Leu Ala Glu Ile Tyr
325 330 335

Val Asn Ser Ser Phe Tyr Lys Glu Thr Lys Ala Glu Leu His Gln Leu
340 345 350

Ser Gly Gly Glu Lys Lys Lys Lys Ile Thr Val Phe Lys Glu Ile Ser
355 360 365

Tyr Thr Thr Ser Phe Cys His Gln Leu Arg Trp Val Ser Lys Arg Ser
370 375 380

Phe Lys Asn Leu Leu Gly Asn Pro Gln Ala Ser Ile Ala Gln Ile Ile
385 390 395 400

Val Thr Val Val Leu Gly Leu Val Ile Gly Ala Ile Tyr Phe Gly Leu
405 410 415

Lys Asn Asp Ser Thr Gly Ile Gln Asn Arg Ala Gly Val Leu Phe Phe
420 425 430

Leu Thr Thr Asn Gln Cys Phe Ser Ser Val Ser Ala Val Glu Leu Phe
435 440 445

Val Val Glu Lys Lys Leu Phe Ile His Glu Tyr Ile Ser Gly Tyr Tyr
450 455 460

Arg Val Ser Ser Tyr Phe Leu Gly Lys Leu Leu Ser Asp Leu Leu Pro
465 470 475 480

Met Arg Met Leu Pro Ser Ile Ile Phe Thr Cys Ile Val Tyr Phe Met
485 490 495

Leu Gly Leu Lys Pro Lys Ala Asp Ala Phe Phe Val Met Met Phe Thr
500 505 510

Leu Met Met Val Ala Tyr Ser Ala Ser Ser Met Ala Leu Ala Ile Ala
515 520 525

Ala Gly Gln Ser Val Val Ser Val Ala Thr Leu Leu Met Thr Ile Cys
530 535 540

Phe Val Phe Met Met Ile Phe Ser Gly Leu Leu Val Asn Leu Thr Thr
545 550 555 560

Ile Ala Ser Trp Leu Ser Trp Leu Gln Tyr Phe Ser Ile Pro Arg Tyr
565 570 575

Gly Phe Thr Ala Leu Gln His Asn Glu Phe Leu Gly Gln Asn Phe Cys
580 585 590

Pro Gly Leu Asn Ala Thr Gly Asn Asn Pro Cys Asn Tyr Ala Thr Cys
595 600 605

Thr Gly Glu Glu Tyr Leu Val Lys Gln Gly Ile Asp Leu Ser Pro Trp
610 615 620

Gly Leu Trp Lys Asn His Val Ala Leu Ala Cys Met Ile Val Ile Phe
625 630 635 640

Leu Thr Ile Ala Tyr Leu Lys Leu Leu Phe Leu Lys Lys Tyr Ser

645

650

655

<210> 3

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 1 forward
primer

<400> 3

gtgccctc aaaggtt

18

<210> 4

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 1 reverse
primer

<400> 4

tccagtcaaa gctgtactct g

21

<210> 5

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 2 forward
primer

<400> 5

atgtattgtc acctagtgtt tg

22

<210> 6

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 2 reverse
primer

<400> 6

aaagtgtgaa gccttgagca ga

22

<210> 7

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 3 forward
primer

<400> 7

aacggagatg tttcacaaga

20

<210> 8

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 3 reverse
primer

<400> 8

tacaataaag ccccaaaaca

20

<210> 9
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Exon 4 forward
primer

<400> 9
gaggaaaaag aatgggagaa

20

<210> 10
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Exon 4 reverse
primer

<400> 10
gtctgcaaag cctgctataa

20

<210> 11
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Exon 5 forward
primer

<400> 11
ttccttcacc tttcttttcc

20

<210> 12

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 5 reverse
primer

<400> 12

cttcataaa actggtcct

20

<210> 13

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 6 forward
primer

<400> 13

gaggtgcttt gtatcaggct

20

<210> 14

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 6 reverse
primer

<400> 14

gatcaggcca gtaggtcaac

20

<210> 15

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 7 forward
primer

<400> 15

cttgtaaata cttgcagatt acctg

25

<210> 16

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 7 reverse
primer

<400> 16

tgttcaagtg acagaataaa tggct

25

<210> 17

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 8 forward
primer

<400> 17

aaagggtaaa attacgtggg

20

<210> 18

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 8 reverse
primer

<400> 18

gcaaacaac tgacgttttc

20

<210> 19

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 9 forward
primer

<400> 19

aatgaagggtg ttagggaagc

20

<210> 20

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 9 reverse
primer

<400> 20

ctggctgaca cttctttcac

20

<210> 21

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 10 forward
primer

<400> 21

tctccccaaa gcacagataa ct

22

<210> 22

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 10 reverse
primer

<400> 22

catttaaaaa taattgggcc aggtg

25

<210> 23

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 11 forward
primer

<400> 23

ctaattacct tccaaagggc

20

<210> 24

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 11 reverse primer

<400> 24

aaaccaggct gctctttact

20

<210> 25

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 12 forward primer

<400> 25

gctgggtatt tttcaaggat

20

<210> 26

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 12 reverse primer

<400> 26

agagagtgca aaatggacag

20

<210> 27

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 13 forward
primer

<400> 27

tgccctgtagc tcttcatctc

20

<210> 28

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 13 reverse
primer

<400> 28

acgagaggga accaaaatag

20

<210> 29

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 14 forward
primer

<400> 29

ctttttggca gctttaaatg atagc

25

<210> 30

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 14 reverse
primer

<400> 30

aatctttctc ctttactagg aggta

25

<210> 31

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 15 forward
primer

<400> 31

tttacttctt ttgtattgga agcca

25

<210> 32

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 15 reverse
primer

<400> 32

tagaggataa atcgattgat aggga

25

<210> 33

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 16 forward primer

<400> 33

atctgaagg gtaattatta aaggc

25

<210> 34

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 16 reverse primer

<400> 34

tgttccagaa atggtgcaag aattc

25

<210> 35

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 1 sense primer

<400> 35

gtgccactc aaaaggtt

18

<210> 36

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 1

antisense primer

<400> 36

caagagtttt taccaaccca

20

<210> 37

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 2 sense
primer

<400> 37

atgtattgtc acctagtgtt tg

22

<210> 38

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 2
antisense primer

<400> 38

gtggcccaat tatttcact

19

<210> 39

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 3 sense
primer

<400> 39

taagagtgg ttgtgcttg

20

<210> 40

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 3
antisense primer

<400> 40

aacatggta actgctacat

20

<210> 41

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 4 sense
primer

<400> 41

atgttttggg gctttattg

19

<210> 42

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 4
antisense primer

<400> 42

tattccagat tctccctgc

19

<210> 43

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 5 sense
primer

<400> 43

caggctttgc agacatcta

19

<210> 44

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 5
antisense primer

<400> 44

attgttatgg aaagcaacca

20

<210> 45

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 6 sense
primer

<400> 45

gaggtgcttt gtatcaggct

20

<210> 46

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 6
antisense primer

<400> 46

caccctcatc acagacatc

19

<210> 47

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 7 sense
primer

<400> 47

ctgtcctaga atctgcattt

20

<210> 48

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 7
antisense primer

<400> 48

agctggtgct acaaaaat

18

<210> 49
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Exon 8 sense
primer

<400> 49
aaagggtaaa attacgtggg

20

<210> 50
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Exon 8
antisense primer

<400> 50
tctggttggt gcttcctact

20

<210> 51
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Exon 9 sense
primer

<400> 51
gttagggaag catccaaga

19

<210> 52
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Exon 9
antisense primer

<400> 52
agggaagctt tccaaaagta

20

<210> 53
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Exon 10 sense
primer

<400> 53
tctccccaaa gcacagataa ct

22

<210> 54
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Exon 10
antisense primer

<400> 54
tggtggtgga tgtctgtagt

20

<210> 55

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 11 sense
primer

<400> 55

ctaattacct tccaaagggc

20

<210> 56

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 11
antisense primer

<400> 56

gctcaggatt ttcttccta

20

<210> 57

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 12 sense
primer

<400> 57

ctggactgag tgttcaggag

20

<210> 58

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 12
antisense primer

<400> 58

agagagtgc aaatggacag

20

<210> 59

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 13 sense
primer

<400> 59

tgccctgtagc tcttcacatc

20

<210> 60

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 13
antisense primer

<400> 60

ataagggcaa agaggaaagt

20

<210> 61

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 14 sense
primer

<400> 61

tttgttcttc ctttaaaacc g

21

<210> 62

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 14
antisense primer

<400> 62

aatctttctc ctttactagg aggta

25

<210> 63

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 15 sense
primer

<400> 63

tttacttctt ttgtattgga agcca

25

<210> 64

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 15
antisense primer

<400> 64

aaaaggccca aaacaataag

20

<210> 65

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 16-1 sense
primer

<400> 65

atctgaaggg gtaattatta aaggc

25

<210> 66

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Exon 16-1
antisense primer

<400> 66

caggagtffc cagaattcaa

20

<210> 67

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 16-2 sense
primer

<400> 67

tggtgttttc tggtcccttg

20

<210> 68

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Exon 16-2
antisense primer

<400> 68

tggtccagaa atggtgcaag aattc

25

[Brief Description of the Drawings]

[Fig.1]

Fig. 1 is a schematic diagram of the ABCG2 polypeptide showing the positions of single nucleotide polymorphisms of the present invention.

[Fig.2]

Fig. 2 is a result of Northern blot analysis by which the amount of the ABCG2 mRNA in various transformed cells was determined.